

(20%) (一) Which of the following statements are true and which is false?

- (1) If $f'(c) = 0$, then $f(x)$ has a maximum or minimum value at $x = c$.
- (2) If $f'(x) = g'(x)$ for all x in an interval I , then $f(x) = g(x)$ on I .
- (3) If $f(x)$ is differentiable on the open interval (a, b) , and c is a point of local maximum for f in (a, b) , then $f'(c) = 0$.
- (4) If $f''(a)$ exists, then f' is continuous at a .
- (5) If $f''(x_0) = 0$, then $(x_0, f(x_0))$ is an inflection point.
- (6) If $\int_a^b f(x)dx = \int_a^b g(x)dx$, then $f(x) = g(x)$.
- (7) If $\lim_{n \rightarrow \infty} a_n = 0$, then $\sum a_n$ converges.
- (8) If $\sum_{n=1}^{\infty} a_n$ converges, then $\lim a_n = 0$.
- (9) If $\sum_{n=1}^{\infty} a_n$ is a series of nonnegative terms, and $a_1 + a_2 + \dots + a_n \leq 5$ for all n , then $\sum a_n$ converges.
- (10) If $\sum_{n=1}^{\infty} a_n$ is a series of positive terms and $\frac{a_n+1}{a_n} < 1$ for all n , then $\sum a_n$ converges.

(60%) (二) Choice.

(1) Find the value of the integral $\int_0^1 x \tan^{-1} dx$.

- (A) $\pi/4$ (B) $\pi - 2$ (C) $\pi/2$ (D) $(\pi - 2)/2$
 (E) $(\pi - 2)/4$ (F) $\pi - 1$ (G) $(\pi - 1)/2$ (H) $(\pi - 1)/4$

(2) Find the value of the integral $\int_0^{1/2} \sqrt{1-x^2} dx$.

- (A) $\frac{\pi}{12} + \frac{\sqrt{3}}{8}$ (B) $\pi/4$ (C) $\pi/12 + \frac{1}{4}$ (D) $\frac{\pi}{12} + \frac{\sqrt{3}}{4}$
 (E) $5/24\pi$ (F) $\frac{1}{12} + \frac{\pi}{4}$ (G) $\frac{\pi}{12} - \frac{\sqrt{3}}{4}$ (H) $\frac{\pi}{12} - \frac{\sqrt{3}}{8}$

(3) Find the value of the integral $\int_0^2 \frac{1}{(x-1)^2} dx$.

- (A) 0 (B) -2 (C) $\frac{3}{2}$ (D) $\ln 5$
 (E) 2 (F) $\frac{\pi}{3}$ (G) $\frac{1}{2}$ (H) diverge.

(4) Find the value of the integral $\int_0^{\pi/4} \sin^{10} x dx$.

- (A) $\frac{1}{2}$ (B) $\frac{\pi}{3}$ (C) 0 (D) 1
 (E) $\frac{63}{512}\pi$ (F) $\frac{63}{216}$ (G) 2π (H) $\frac{63}{108}\pi$

(5) Find the value of the integral $\int_0^{\sqrt{2}} \frac{1}{\sqrt{x^2 + 1}} dx$.

- (A) $\sqrt{3}/4$ (B) 2 (C) $\ln 4$ (D) $\ln(\sqrt{5}/4)$
 (E) $\sqrt{5}/4$ (F) $\ln 2$ (G) $\sqrt{2}$ (H) $\ln(\sqrt{3}/4)$

(6) Find the shortest distance from the point $(1, 4)$ to a point on the parabola $y^2 = 2x$.

- (A) 1 (B) $\sqrt{2}$ (C) $\sqrt{3}$ (D) 2
 (E) $\sqrt{5}$ (F) $\sqrt{6}$ (G) $\sqrt{7}$ (H) $2\sqrt{2}$

(7) Find the area of the largest rectangle that can be inscribed in the ellipse

$$x^2 + \frac{y^2}{4} = 1.$$

- (A) 1 (B) 2 (C) 3 (D) 4
 (E) 10 (F) 6 (G) 7 (H) 8

(8) Find the value of the limit $\lim_{n \rightarrow \infty} \sum_{i=1}^n \frac{1}{n} \left[\left(\frac{i}{n} \right)^3 + 1 \right]$.

- (A) 0 (B) $\frac{1}{4}$ (C) $\frac{1}{2}$ (D) $\frac{1}{3}$
 (E) $\frac{3}{4}$ (F) $\frac{2}{3}$ (G) 1 (H) $\frac{5}{4}$

(9) Let $f(x) = \int_{\sqrt{x}}^x \cos t dt$. Find the value of $f'(1)$.

(A) 0 (B) $\cos 1$ (C) $2\cos 1$ (D) $\frac{3}{2}\cos 1$

(E) $\sin 1$ (F) $2\sin 1$ (G) $\frac{3\sin 1}{2}$ (H) $\frac{3}{2}\sin 1$

(10) Find the average value of $f(x) = 4\sqrt{x+1}$ on $[0, 15]$.

(A) 1 (B) 56 (C) $\frac{56}{3}$ (D) 28

(E) $\frac{\pi}{2}$ (F) $\frac{56}{5}$ (G) 16 (H) $\frac{16}{15}$

(11) Which of the following series are conditionally convergent?

i) $\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{\sqrt{n}}$ ii) $\sum_{n=1}^{\infty} \frac{n^n}{n!}$ iii) $\sum_{n=1}^{\infty} \frac{\sin n\pi}{\pi^n}$

(A) none (B) i (C) ii (D) iii
(E) i, ii (F) i, iii (G) ii, iii (H) i, ii, iii

(12) Find the value of the series $\sum_{n=1}^{\infty} \frac{1}{n2^n}$.

(A) $\frac{1}{2}$ (B) $\sqrt{2}$ (C) 2 (D) $\ln 2$

(E) $\frac{1}{\ln 2}$ (F) $\sqrt{\ln 2}$ (G) 4 (H) $2\ln 2$

(13) Find the sum of the series $\sum_{n=1}^{\infty} nx^n$.

(A) $\frac{1}{1-x}$ (B) $\frac{1}{(1-x)^2}$ (C) $\frac{x}{(1-x)^2}$ (D) $\frac{x}{1-x}$

(E) $\sqrt{1-x}$ (F) $\frac{1}{1+x}$ (G) $\frac{1}{1+x^2}$ (H) $\frac{x}{1+x^2}$

(14) Find the value of $\lim_{n \rightarrow \infty} \left(1 - \frac{3}{n}\right)^{\frac{n}{2}}$.

(A) 1 (B) e (C) e^3 (D) e^{-3}

- (E) $e^{\frac{1}{2}}$ (F) $e^{-\frac{1}{2}}$ (G) $e^{\frac{1}{2}}$ (H) $e^{-\frac{1}{2}}$

(15) Let $f(x) = (\sin x)^x$. Find the value of $f'(\frac{\pi}{4})$.

- (A) $\frac{1}{2}$ (B) 1 (C) 0 (D) -1
(E) 2 (F) $\sqrt{2}$ (G) $-\sqrt{2}$ (H) $-\frac{1}{2}$

(5%) (三) Use polar coordinates to find the volume below the paraboloid $z = 4 - x^2 - y^2$ and above the xy -plane.

(15%) (四) The Gamma function, defined by $P(\alpha) = \int_0^\infty x^{\alpha-1} e^{-x} dx$, $\alpha > 0$ has many

application in both pure and applied mathematics. Show the following

(a) The integral that defines $P(\alpha)$ converges, if $\alpha > 0$

(b) $P(\alpha + 1) = \alpha P(\alpha)$, $\alpha > 0$

(c) Find the value of $P(5) = \int_0^\infty x^4 e^{-x} dx$

(d) Find the value of $P(\frac{1}{2})$

(e) Find the value of the integral $\int_0^\infty x^2 e^{-\frac{x}{2}} dx$.