

本試題是否可以使用計算機： 可使用， 不可使用（請命題老師勾選）

考試日期：0301，節次：3

1. (15%) Let  $\mathbf{A}$  be an  $n \times n$  symmetric matrix. If  $\lambda_1$  and  $\lambda_2$  are distinct eigenvalues of  $\mathbf{A}$ , show that their corresponding eigenvectors  $\mathbf{x}_1$  and  $\mathbf{x}_2$  are orthogonal.

2. (30%) Consider line integrals  $\int_c \mathbf{F} \cdot d\mathbf{r} = \int_c F_1 dx + F_2 dy + F_3 dz$ , where  $\mathbf{F} = (F_1, F_2, F_3)$ ,  $\mathbf{r} = (x, y, z)$  are vectors, prove that this line integral is path independent if and only if

(a)  $\mathbf{F} = \text{grad } f = \nabla f$

or (b)  $\oint_c \mathbf{F} \cdot d\mathbf{r} = 0$  (integration around closed curves  $c$  always gives 0)

or (c)  $\nabla \times \mathbf{F} = 0$  provided the region enclosed by curve  $c$  is simply connected.

Note: The expression " $\cdot$ " and " $\times$ " represents the dot and cross product, respectively.

3. (20%) In an undamped mass-spring system, resonance occurs if the frequency of the driving force equals the natural frequency of the system and the model can be written as

$$y'' + \omega_0^2 y = K \sin \omega_0 t \quad (1),$$

where  $y(0) = y'(0) = 0$  and  $K$  is constant.

Solve equation (1) with given initial conditions using Laplace transform.

Hint: Use the convolution integral theorem:  $\mathcal{L}^{-1}(F(s)G(s)) = f * g$

4. (15%) Evaluate  $\int_0^{2\pi} \frac{1}{(2+\cos \theta)^2} d\theta$  using contour integral

5. (20%) Solve the non-homogeneous diffusion problem

$$u_t - c^2 u_{xx} = e^{-\alpha x} \quad 0 < x < L, \text{ where } c \text{ and } \alpha \text{ are constant.}$$

$$\text{BC's: } u(0, t) = u(L, t) = 0$$

$$\text{IC: } u(x, 0) = f(x)$$