

本試題是否可以使用計算機：可使用，不可使用（請命題老師勾選）

考試日期：0301，節次：3

1. Please prove the following addition formulas for cosine and sine: (10%)

$$\begin{cases} \sin(\alpha + \beta) = \sin \alpha \cos \beta + \cos \alpha \sin \beta \\ \sin(\alpha - \beta) = \sin \alpha \cos \beta - \cos \alpha \sin \beta \\ \cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta \\ \cos(\alpha - \beta) = \cos \alpha \cos \beta + \sin \alpha \sin \beta \end{cases}$$

(Hint: You can use the linear transformation $y=Ax$ with a rotation matrix A to derive them.)

2. The derivative of a continuous function $f(x)$ is defined by $f'(x) = \frac{df}{dx}$

$$= \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x}, \forall x. \text{ Please prove the derivatives of } \sin(x) \text{ and } \cos(x) \text{ are}$$

$$\sin'(x) = \cos(x) \text{ and } \cos'(x) = -\sin(x), \text{ respectively. (10\%)}$$

3. Please find the curve through the point (2,4) in the xy -plane having the slope $-y/x$ at each point of the curve. (10%)

4. Please find the particular solution of the initial value problem $y' = -2xy$ for an unknown function $y(x)$ with the initial condition $y(0)=1$. (10%)

5. A real square matrix $A = \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix}$ is given.

(5a) Please determine the eigenvalues and eigenvectors of A . (10%)

(5b) This matrix A is orthogonal. All orthogonal matrices have some special properties, e.g. ---

① orthogonality : please state the definition of an orthogonal matrix and verify that A is an orthogonal matrix. (5%)

(背面仍有題目,請繼續作答)

本試題是否可以使用計算機： 可使用， 不可使用（請命題老師勾選）

考試日期：0301，節次：3

② invariance of inner product: An orthogonal transformation preserves the value of the inner product of vectors. What is an orthogonal transformation? Please prove it and verify that it holds for this matrix A. (8%)

③ determinant of an orthogonal matrix: please compute the determinant of A and prove that the determinant of an orthogonal matrix has the value +1 or -1. (8%)

6. Let R be a **closed bounded region** in the xy -plane whose boundary C consists of finitely many smooth curves. Let $F_1(x, y)$ and $F_2(x, y)$ be functions that are continuous and have continuous partial derivatives $\partial F_1/\partial y$ and $\partial F_2/\partial x$ everywhere in some domain

containing R . Then $\iint_R \left(\frac{\partial F_2}{\partial x} - \frac{\partial F_1}{\partial y} \right) dx dy = \oint_C (F_1 dx + F_2 dy)$. This is the famous **Green's**

theorem in the plane.

(6a) Please verify it for $F_1 = 6y$, $F_2 = 2x + 2$ and C the circle $x^2 + y^2 = 1$. (10%)

(6b) After derivation, we have $A = \frac{1}{2} \oint_C (x dy - y dx)$. This formula expresses the area of R in terms of a line integral over the boundary. It has various applications, e.g. the theory of certain **planimeters** (instruments for measuring area) is based on

it. Please derive this formula $A = \frac{1}{2} \oint_C (x dy - y dx)$. (12%)

(6c) Please prove that an ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ has the area $A = \pi ab$. (7%)