※ 考生請注意：本試題不可使用計算機。　請於答案卷(卡)作答，於本試題紙上作答者，不予計分。

選擇題（選出正確答案，單選題，每題四分）

1. If A and B are independent events with P(A) = 0.65 and P(A ∩ B)= 0.26, then, P(B) =
   a. 0.400
   b. 0.169
   c. 0.390
   d. 0.650

2. A six-sided die is tossed 3 times.　The probability of observing three ones in a row is
   a. 1/3
   b. 1/6
   c. 1/27
   d. 1/216

3. A simple random sample of 100 observations was taken from a large population.　The sample mean and the standard deviation were determined to be 80 and 12 respectively.　The standard error of the mean is
   a. 1.20
   b. 0.12
   c. 8.00
   d. 0.80

4. A random sample of 16 statistics examinations was taken.　The average score, in the sample, was 76.2 with a variance of 144.　The 99% confidence interval for the true average examination score is
   a. 49.677 to 102.723
   b. 73.99 to 78.41
   c. 67.359 to 85.041
   d. 67.437 to 84.963

5. An interval estimate is a range of values used to estimate
   a. the shape of the population's distribution
   b. the sampling distribution
   c. a sample statistic
   d. a population parameter

Exhibit AA

Salary information for a random sample of male and female employees of a large company is shown below.

	Male	Female
Sample Size	64	36
Sample Mean Salary (in $1,000)	44	41
Sample Variance	128	72

6. Refer to Exhibit AA.　The standard error for the difference between the two means is
   a. 4
   b. 7.46
   c. 4.24
   d. 2.0

7. Refer to Exhibit AA.　The 95% confidence interval for the difference between the means of the two populations is
   a. 0 to 6.92
   b. -2 to 2
   c. -1.96 to 1.96
   d. -0.92 to 6.92

※ 考生請注意：本試題不可使用計算機。　請於答案卷(卡)作答，於本試題紙上作答者，不予計分。

8.　　The producer of a certain medicine claims that their bottling equipment is very accurate and that the standard deviation of all their filled bottles is 0.1 ounce or less. A sample of 20 bottles showed a standard deviation of 0.11. The test statistic to test the claim is
  a.　400
  b.　22.99
  c.　4.85
  d.　20

Exhibit BB
The table below gives beverage preferences for random samples of teens and adults.

	Teens	Adults	Total
Coffee	50	200	250
Tea	100	150	250
Soft Drink	200	200	400
Other	50	50	100
	400	600	1,000

We are asked to test for independence between age (i.e., adult and teen) and drink preferences.

9.　　Refer to Exhibit BB. With a .05 level of significance, the critical value for the test is
  a.　1.645
  b.　7.815
  c.　14.067
  d.　15.507

10.　　Refer to Exhibit BB. The expected number of adults who prefer coffee is
  a.　0.25
  b.　0.33
  c.　150
  d.　200

Exhibit CC

  $n = 16$　　$\overline{X} = 75.76$　$s = 8.246$　　$H_0:\ \mu \geq 80$
  $H_a:\ \mu < 80$

11.　　Refer to Exhibit CC. The standardized test statistic equals
  a.　-2.12
  b.　-0.53
  c.　0.53
  d.　0.483

12.　　Refer to Exhibit CC. The p-value is equal to
  a.　-0.025
  b.　0.05
  c.　0.525
  d.　0.025

※ 考生請注意：本試題不可使用計算機。　請於答案卷(卡)作答，於本試題紙上作答者，不予計分。

13. If a hypothesis is rejected at 95% confidence, it
   a. will always be accepted at 90% confidence
   b. will always be rejected at 90% confidence
   c. will sometimes be rejected at 90% confidence
   d. None of these alternatives is correct.

Exhibit DD
You are given the following information about y and x.

y Dependent Variable	x Independent Variable
5	15
7	12
9	10
11	7

14. Refer to Exhibit DD. The least squares estimate of b_0 equals
   a. -0.7647
   b. -1.3
   c. 164.1176
   d. 16.41176

15. Refer to Exhibit DD. The coefficient of determination equals
   a. -0.99705
   b. -0.9941
   c. 0.9941
   d. 0.99705

16. If the coefficient of correlation is a positive value, then
   a. the intercept must also be positive
   b. the coefficient of determination can be either negative or positive, depending on the value of the slope
   c. the regression equation could have either a positive or a negative slope
   d. the slope of the line must be positive

17. In multiple regression analysis, the general linear model
   a. can not be used to accommodate curvilinear relationships between dependent variables and independent variables
   b. can be used to accommodate curvilinear relationships between the independent variables and dependent variable
   c. must contain more than 2 independent variables
   d. None of these alternatives is correct.

18. In an analysis of variance problem involving 3 treatments and 10 observations per treatment, SSE = 399.6. The MSE for this situation is
   a. 133.2
   b. 13.32
   c. 14.8
   d. 30.0

※ 考生請注意：本試題不可使用計算機。　請於答案卷(卡)作答，於本試題紙上作答者，不予計分。

19. When an analysis of variance is performed on samples drawn from K populations, the mean square between treatments (MSTR) is
   a. $SSTR/n_T$
   b. $SSTR/(n_T - 1)$
   c. $SSTR/K$
   d. $SSTR/(K - 1)$
   e. None of these alternatives is correct.

Exhibit EE

   $f(x) = (1/10)\ e^{-x/10}\quad x \geq 0$

20. Refer to Exhibit EE.　The mean of x is
   a. 0.10
   b. 10
   c. 100
   d. 1,000

21. Refer to Exhibit EE.　The probability that x is between 3 and 6 is
   a. 0.4512
   b. 0.1920
   c. 0.2592
   d. 0.6065

22. In a regression model involving more than one independent variable, which of the following tests must be used in order to determine if the relationship between the dependent variable and the set of independent variables is significant?
   a. t test
   b. F test
   c. Either a t test or a chi-square test can be used.
   d. chi-square test

23. In simple linear regression analysis, which of the following is not true?
   a. The F test and the t test yield the same results.
   b. The F test and the t test may or may not yield the same results.
   c. The relationship between X and Y is represented by means of a straight line.
   d. The value of $F = t^2$.

※ 考生請注意：本試題不可使用計算機。　請於答案卷(卡)作答，於本試題紙上作答者，不予計分。

Exhibit FF

You want to test whether or not the following sample of 30 observations follows a normal distribution. The mean of the sample equals 11.83, and the standard deviation equals 4.53. The number of intervals or categories used to test the hypothesis for this problem is 6.

| 2	3	5	5	7	8	8	9	9	10
11	11	12	12	12	12	13	13	13	14
15	15	15	16	16	17	17	18	18	19

24. Refer to Exhibit FF. The calculated value for the test statistic equals
   a. 0
   b. 1.67
   c. 2
   d. 6

25. Refer to Exhibit FF. The hypothesis is to be tested at the 5% level of significance. The critical value from the table equals
   a. 1.645
   b. 1.96
   c. 7.815
   d. 12.592

標準常態分配

Z

累積機率

表中的數值代表對應的 z 值以左，標準常態曲線下方的面積。例如：z = 1.25，累積機率爲 0.8944。

z	0.00	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09
0.0	0.5000	0.5040	0.5080	0.5120	0.5160	0.5199	0.5239	0.5279	0.5319	0.5359
0.1	0.5398	0.5438	0.5478	0.5517	0.5557	0.5596	0.5636	0.5675	0.5714	0.5753
0.2	0.5793	0.5832	0.5871	0.5910	0.5948	0.5987	0.6026	0.6064	0.6103	0.6141
0.3	0.6179	0.6217	0.6255	0.6293	0.6331	0.6368	0.6406	0.6443	0.6480	0.6517
0.4	0.6554	0.6591	0.6628	0.6664	0.6700	0.6736	0.6772	0.6808	0.6844	0.6879
0.5	0.6915	0.6950	0.6985	0.7019	0.7054	0.7088	0.7123	0.7157	0.7190	0.7224
0.6	0.7257	0.7291	0.7324	0.7357	0.7389	0.7422	0.7454	0.7486	0.7517	0.7549
0.7	0.7580	0.7611	0.7642	0.7673	0.7704	0.7734	0.7764	0.7794	0.7823	0.7852
0.8	0.7881	0.7910	0.7939	0.7967	0.7995	0.8023	0.8051	0.8078	0.8106	0.8133
0.9	0.8159	0.8186	0.8212	0.8238	0.8264	0.8289	0.8315	0.8340	0.8365	0.8389
1.0	0.8413	0.8438	0.8461	0.8485	0.8508	0.8531	0.8554	0.8577	0.8599	0.8621
1.1	0.8643	0.8665	0.8686	0.8708	0.8729	0.8749	0.8770	0.8790	0.8810	0.8830
1.2	0.8849	0.8869	0.8888	0.8907	0.8925	0.8944	0.8962	0.8980	0.8997	0.9015
1.3	0.9032	0.9049	0.9066	0.9082	0.9099	0.9115	0.9131	0.9147	0.9162	0.9177
1.4	0.9192	0.9207	0.9222	0.9236	0.9251	0.9265	0.9279	0.9292	0.9306	0.9319
1.5	0.9332	0.9345	0.9357	0.9370	0.9382	0.9394	0.9406	0.9418	0.9429	0.9441
1.6	0.9452	0.9463	0.9474	0.9484	0.9495	0.9505	0.9515	0.9525	0.9535	0.9545
1.7	0.9554	0.9564	0.9573	0.9582	0.9591	0.9599	0.9608	0.9616	0.9625	0.9633
1.8	0.9641	0.9649	0.9656	0.9664	0.9671	0.9678	0.9686	0.9693	0.9699	0.9706
1.9	0.9713	0.9719	0.9726	0.9732	0.9738	0.9744	0.9750	0.9756	0.9761	0.9767
2.0	0.9772	0.9778	0.9783	0.9788	0.9793	0.9798	0.9803	0.9808	0.9812	0.9817
2.1	0.9821	0.9826	0.9830	0.9834	0.9838	0.9842	0.9846	0.9850	0.9854	0.9857
2.2	0.9861	0.9864	0.9868	0.9871	0.9875	0.9878	0.9881	0.9884	0.9887	0.9890
2.3	0.9893	0.9896	0.9898	0.9901	0.9904	0.9906	0.9909	0.9911	0.9913	0.9913
2.4	0.9918	0.9920	0.9922	0.9925	0.9927	0.9929	0.9931	0.9932	0.9934	0.9936
2.5	0.9938	0.9940	0.9941	0.9943	0.9945	0.9946	0.9948	0.9949	0.9951	0.9952
2.6	0.9953	0.9955	0.9956	0.9957	0.9959	0.9960	0.9961	0.9962	0.9963	0.9964
2.7	0.9965	0.9966	0.9967	0.9968	0.9969	0.9970	0.9971	0.9972	0.9973	0.9974
2.8	0.9974	0.9975	0.9976	0.9977	0.9977	0.9978	0.9979	0.9979	0.9980	0.9981
2.9	0.9981	0.9982	0.9982	0.9983	0.9984	0.9984	0.9985	0.9985	0.9986	0.9986
3.0	0.9986	0.9987	0.9987	0.9988	0.9988	0.9989	0.9989	0.9989	0.9990	0.9990

t分配之臨界值

γ	0.10	0.05	0.025	0.01	0.005
1	3.078	6.314	12.706	31.821	63.657
2	1.886	2.920	4.303	6.965	9.925
3	1.638	2.353	3.182	4.541	5.841
4	1.533	2.132	2.776	3.747	4.604
5	1.476	2.015	2.571	3.365	4.032
6	1.440	1.943	2.447	3.143	3.707
7	1.415	1.895	2.365	2.998	3.499
8	1.397	1.860	2.306	2.896	3.355
9	1.383	1.833	2.262	2.821	3.250
10	1.372	1.812	2.228	2.764	3.169
11	1.363	1.796	2.201	2.718	3.106
12	1.356	1.782	2.179	2.681	3.055
13	1.350	1.771	2.160	2.650	3.012
14	1.345	1.761	2.145	2.624	2.977
15	1.341	1.753	2.131	2.602	2.947
16	1.337	1.746	2.120	2.583	2.921
17	1.333	1.740	2.110	2.567	2.898
18	1.330	1.734	2.101	2.552	2.878
19	1.328	1.729	2.093	2.539	2.861
20	1.325	1.725	2.086	2.528	2.845
21	1.323	1.721	2.080	2.518	2.831
22	1.321	1.717	2.074	2.508	2.819
23	1.319	1.714	2.069	2.500	2.807
24	1.318	1.711	2.064	2.492	2.797
25	1.316	1.708	2.060	2.485	2.787
26	1.315	1.706	2.056	2.479	2.779
27	1.314	1.703	2.052	2.473	2.771
28	1.313	1.701	2.048	2.467	2.763
29	1.311	1.699	2.045	2.462	2.756
inf.	1.282	1.645	1.960	2.326	2.576

卡方

自由度	機率 $1-\alpha$							
	.005	.010	.025	.050	.950	.975	.990	.995
1				.004	3.84	5.02	6.63	7.88
2	.01	.02	.05	.10	5.99	7.38	9.21	10.60
3	.07	.11	.22	.35	7.81	9.35	11.34	12.84
4	.21	.30	.48	.71	9.49	11.14	13.28	14.86
5	.41	.55	.83	1.15	11.07	12.83	15.09	16.75
6	.68	.87	1.24	1.64	12.59	14.45	16.81	18.55
7	.99	1.24	1.69	2.17	14.07	16.01	18.48	20.28
8	1.34	1.65	2.18	2.73	15.51	17.53	20.09	21.96
9	1.73	2.09	2.70	3.33	16.92	19.02	21.67	23.59
10	2.16	2.56	3.25	3.94	18.31	20.48	23.21	25.19
11	2.60	3.05	3.82	4.57	19.68	21.92	24.72	26.76
12	3.07	3.57	4.40	5.23	21.03	23.34	26.22	28.30
13	3.57	4.11	5.01	5.89	22.36	24.74	27.69	29.82
14	4.07	4.66	5.63	6.57	23.68	26.12	29.14	31.32
15	4.60	5.23	6.26	7.26	25.00	27.49	30.58	32.80
16	5.14	5.81	6.91	7.96	26.30	28.85	32.00	34.27
17	5.70	6.41	7.56	8.67	27.59	30.19	33.41	35.72
18	6.26	7.01	8.23	9.39	28.87	31.53	34.81	37.16
19	6.84	7.63	8.91	10.12	30.14	32.85	36.19	38.58
20	7.43	8.26	9.59	10.85	31.41	34.17	37.57	40.00
21	8.03	8.90	10.28	11.59	32.67	35.48	38.93	41.40
22	8.64	9.54	10.98	12.34	33.92	36.78	40.29	42.80
23	9.26	10.20	11.69	13.09	35.17	38.08	41.64	44.18
24	9.89	10.86	12.40	13.85	36.42	39.36	42.98	45.56
25	10.52	11.52	13.12	14.61	37.65	40.65	44.31	46.93
26	11.16	12.20	13.84	15.38	38.89	41.92	45.64	48.29
27	11.81	12.88	14.57	16.15	40.11	43.19	46.96	49.64
28	12.46	13.56	15.31	16.93	41.34	44.46	48.28	50.99
29	13.12	14.26	16.05	17.71	42.56	45.72	49.59	52.34
30	13.79	14.95	16.79	18.49	43.77	46.98	50.89	53.67

$1-\alpha = 0.95$

ν_2 \ ν_1	1	2	3	4	5	6	7	8	9
1	161.45	199.50	215.71	224.58	230.16	233.99	236.77	238.88	240.54
2	18.513	19.000	19.164	19.247	19.296	19.330	19.353	19.371	19.385
3	10.128	9.5521	9.2766	9.1172	9.0135	8.9406	8.8868	8.8452	8.8123
4	7.7086	6.9443	6.5914	6.3883	6.2560	6.1631	6.0942	6.0410	5.9988
5	6.6079	5.7861	5.4095	5.1922	5.0503	4.9503	4.8759	4.8183	4.7725
6	5.9874	5.1433	4.7571	4.5337	4.3874	4.2839	4.2066	4.1468	4.0990
7	5.5914	4.7374	4.3468	4.1203	3.9715	3.8660	3.7870	3.7257	3.6767
8	5.3177	4.4590	4.0662	3.8378	3.6875	3.5806	3.5005	3.4381	3.3881
9	5.1174	4.2565	3.8626	3.6331	3.4817	3.3738	3.2927	3.2296	3.1789
10	4.9646	4.1028	3.7083	3.4780	3.3258	3.2172	3.1355	3.0717	3.0204
11	4.8443	3.9823	3.5874	3.3567	3.2039	3.0946	3.0123	2.9480	2.8962
12	4.7472	3.8853	3.4903	3.2592	3.1059	2.9961	2.9134	2.8486	2.7964
13	4.6672	3.8056	3.4105	3.1791	3.0254	2.9153	2.8321	2.7669	2.7144
14	4.6001	3.7389	3.3439	3.1122	2.9582	2.8477	2.7642	2.6987	2.6458
15	4.5431	3.6823	3.2874	3.0556	2.9013	2.7905	2.7066	2.6408	2.5876
16	4.4940	3.6337	3.2389	3.0069	2.8524	2.7413	2.6572	2.5911	2.5377
17	4.4513	3.5915	3.1968	2.9647	2.8100	2.6987	2.6143	2.5480	2.4943
18	4.4139	3.5546	3.1599	2.9277	2.7729	2.6613	2.5767	2.5102	2.4563
19	4.3808	3.5219	3.1274	2.8951	2.7401	2.6283	2.5435	2.4768	2.4227
20	4.3513	3.4928	3.0984	2.8661	2.7109	2.5990	2.5140	2.4471	2.3928
21	4.3248	3.4668	3.0725	2.8401	2.6848	2.5757	2.4876	2.4205	2.3661
22	4.3009	3.4434	3.0491	2.8167	2.6613	2.5491	2.4638	2.3965	2.3419
23	4.2793	3.4221	3.0280	2.7955	2.6400	2.5277	2.4422	2.3748	2.3201
24	4.2597	3.4028	3.0088	2.7763	2.6207	2.5082	2.4226	2.3551	2.3002
25	4.2417	3.3852	2.9912	2.7587	2.6030	2.4904	2.4047	2.3371	2.2821
26	4.2252	3.3690	2.9751	2.7426	2.5868	2.4741	2.3883	2.3205	2.2655
27	4.2100	3.3541	2.9604	2.7278	2.5719	2.4591	2.3732	2.3053	2.2501
28	4.1960	3.3404	2.9467	2.7141	2.5581	2.4453	2.3593	2.2913	2.2360
29	4.1830	3.3277	2.9340	2.7014	2.5454	2.4324	2.3463	2.2782	2.2229
30	4.1709	3.3158	2.9223	2.6896	2.5336	2.4205	2.3343	2.2662	2.2107
40	4.0848	3.2317	2.8387	2.6060	2.4495	2.3359	2.2490	2.1802	2.1240
60	4.0012	3.1504	2.7581	2.5252	2.3683	2.2540	2.1665	2.0970	2.0401
120	3.9201	3.0718	2.6802	2.4472	2.2900	2.1750	2.0867	2.0164	1.9588
∞	3.8415	2.9957	2.6049	2.3719	2.2141	2.0986	2.0096	1.9384	1.8799