

本試題是否可以使用計算機： 可使用， 不可使用 (請命題老師勾選)

Answer all the questions in the paper provided, and show your work. This exam is worth 100 points. Points are shown at the beginning of each question. The various distribution tables are attached to the end of the exam. Pace yourself carefully.

1. [20 points] Suppose that X and Y are continuous random variables with the joint probability density function

$$f(x, y) = \begin{cases} k(1-x)(2-y) & \text{for } 0 \leq x \leq 1, 0 \leq y \leq 2 \\ 0 & \text{otherwise} \end{cases}$$

- (a) Find k .
(b) Find the expected values of X and Y , i.e., $E(X)$ and $E(Y)$.
2. [20 points] Two independent random variables X_1 and X_2 are both normally distributed with same mean but different variance. In particular, they are distributed as $X_1 \sim N(\mu, \sigma_1^2)$, $X_2 \sim N(\mu, \sigma_2^2)$.

- (a) The following estimator for μ is proposed, $\hat{\mu}_A = \frac{X_1 + X_2}{2}$. Is this an unbiased estimator of μ ? What is the variance of this estimator?
(b) Another estimator for μ is also proposed, $\hat{\mu}_B = \frac{1}{3}X_1 + \frac{2}{3}X_2$. Which estimator would you prefer? Why?

3. [20 points] Let X and Y be random variables, let μ_X and σ_X^2 be the mean and variance of X , let σ_{XY} be the covariance between X and Y , let ρ_{XY} be the correlation coefficient between X and Y (and so forth for the other variables), and let a and b be constants. Use the definitions of the mean, variance and/or correlation to answer the following questions:

(a) Show that $\text{var}(aX + bY) = a^2\sigma_X^2 + 2ab\sigma_{XY} + b^2\sigma_Y^2$.

(b) Show that $E(Y^2) = \sigma_Y^2 + \mu_Y^2$.

(背面仍有題目,請繼續作答)

(c) Use the result in part (a) to prove the correlation inequality; that is, $|\rho_{XY}| \leq 1$.

4. [20 points] Suppose that a light bulb manufacturing plant produces bulbs with a mean life of 2000 hours and a standard deviation of 200 hours. An inventor claims to have developed an improved process that produces bulbs with a longer mean life and the same standard deviation. The plant manager randomly selects 100 bulbs produced by the process. She says that she will believe the inventor's claim if the sample mean life of the bulbs is greater than 2100 hours, otherwise she will conclude that the new process is no better than the old process. Let μ denote the mean of the new process. Consider the null and alternative hypothesis $H_0: \mu = 2000$ vs. $H_1: \mu > 2000$.

(a) Suppose that the new process is in fact better and has a mean bulb life of 2150 hours. What is the power of the plant manager's testing procedure?

(b) What testing procedure should the plant manager use if she wants the size of her test to be 5%?

5. [20 points] To investigate possible gender discrimination in a firm, a sample of 100 men and 64 women with similar job descriptions are selected at random. A summary of the resulting monthly salaries are:

	Average Salary (\bar{Y})	Standard Deviation (s_Y)	n
Men	\$3100	\$200	100
Women	\$2900	\$320	64

(a) What do these data suggest about wage differences in the firm? Do they represent statistically significant evidence that wages of men and women are different?

(b) Do these data suggest that the firm is guilty of gender discrimination in its compensation policies? Explain concisely.

Table 1 Area Under the Standard Normal Distribution

	0.00	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09
0.0	0.0000	0.0040	0.0080	0.0120	0.0160	0.0199	0.0239	0.0279	0.0319	0.0359
0.1	0.0398	0.0438	0.0478	0.0517	0.0557	0.0596	0.0636	0.0675	0.0714	0.0753
0.2	0.0793	0.0832	0.0871	0.0910	0.0948	0.0987	0.1026	0.1064	0.1103	0.1141
0.3	0.1179	0.1217	0.1255	0.1293	0.1331	0.1368	0.1406	0.1443	0.1480	0.1517
0.4	0.1554	0.1591	0.1628	0.1664	0.1700	0.1736	0.1772	0.1808	0.1844	0.1879
0.5	0.1915	0.1950	0.1985	0.2019	0.2054	0.2088	0.2123	0.2157	0.2190	0.2224
0.6	0.2257	0.2291	0.2324	0.2357	0.2389	0.2422	0.2454	0.2486	0.2517	0.2549
0.7	0.2580	0.2611	0.2642	0.2673	0.2704	0.2734	0.2764	0.2794	0.2823	0.2852
0.8	0.2881	0.2910	0.2939	0.2967	0.2995	0.3023	0.3051	0.3079	0.3106	0.3133
0.9	0.3159	0.3186	0.3212	0.3238	0.3264	0.3289	0.3315	0.3340	0.3365	0.3389
1.0	0.3413	0.3438	0.3461	0.3485	0.3508	0.3531	0.3554	0.3577	0.3599	0.3621
1.1	0.3643	0.3665	0.3686	0.3708	0.3729	0.3749	0.3770	0.3790	0.3810	0.3830
1.2	0.3849	0.3869	0.3888	0.3907	0.3925	0.3944	0.3962	0.3980	0.3997	0.4015
1.3	0.4032	0.4049	0.4066	0.4082	0.4099	0.4115	0.4131	0.4147	0.4162	0.4177
1.4	0.4192	0.4207	0.4222	0.4236	0.4251	0.4265	0.4279	0.4292	0.4306	0.4319
1.5	0.4332	0.4345	0.4357	0.4370	0.4382	0.4394	0.4406	0.4418	0.4429	0.4441
1.6	0.4452	0.4463	0.4474	0.4484	0.4495	0.4505	0.4515	0.4525	0.4535	0.4545
1.7	0.4554	0.4564	0.4573	0.4582	0.4591	0.4599	0.4608	0.4616	0.4625	0.4633
1.8	0.4641	0.4649	0.4656	0.4664	0.4671	0.4678	0.4686	0.4693	0.4699	0.4706
1.9	0.4713	0.4719	0.4726	0.4732	0.4738	0.4744	0.4750	0.4756	0.4761	0.4767
2.0	0.4773	0.4778	0.4783	0.4788	0.4793	0.4798	0.4803	0.4808	0.4812	0.4817
2.1	0.4821	0.4826	0.4830	0.4834	0.4838	0.4842	0.4846	0.4850	0.4854	0.4857
2.2	0.4861	0.4864	0.4868	0.4871	0.4875	0.4878	0.4881	0.4884	0.4887	0.4890
2.3	0.4893	0.4896	0.4898	0.4901	0.4904	0.4906	0.4909	0.4911	0.4913	0.4916
2.4	0.4918	0.4920	0.4922	0.4925	0.4927	0.4929	0.4931	0.4932	0.4934	0.4936
2.5	0.4938	0.4940	0.4941	0.4943	0.4945	0.4946	0.4948	0.4949	0.4951	0.4952
2.6	0.4953	0.4955	0.4956	0.4957	0.4959	0.4960	0.4961	0.4962	0.4963	0.4964
2.7	0.4965	0.4966	0.4967	0.4968	0.4969	0.4970	0.4971	0.4972	0.4973	0.4974
2.8	0.4974	0.4975	0.4976	0.4977	0.4977	0.4978	0.4979	0.4979	0.4980	0.4981
2.9	0.4981	0.4982	0.4983	0.4983	0.4984	0.4984	0.4985	0.4985	0.4986	0.4986
3.0	0.4987	0.4987	0.4987	0.4988	0.4988	0.4989	0.4989	0.4989	0.4990	0.4990

Source: This table was generated using the SAS® function PROBNORM.

Table 2 Right-Tail Critical Values for the *t*-distribution

DF	$\alpha = .10$	$\alpha = .05$	$\alpha = .025$	$\alpha = .01$	$\alpha = .005$
1	3.078	6.314	12.706	31.821	63.657
2	1.886	2.920	4.303	6.965	9.925
3	1.638	2.353	3.182	4.541	5.841
4	1.533	2.132	2.776	3.747	4.604
5	1.476	2.015	2.571	3.365	4.032
6	1.440	1.943	2.447	3.143	3.707
7	1.415	1.895	2.365	2.998	3.499
8	1.397	1.860	2.306	2.896	3.355
9	1.383	1.853	2.262	2.821	3.250
10	1.372	1.812	2.228	2.764	3.169
11	1.363	1.796	2.201	2.718	3.106
12	1.356	1.782	2.179	2.681	3.055
13	1.350	1.771	2.160	2.650	3.012
14	1.345	1.761	2.145	2.624	2.977
15	1.341	1.753	2.131	2.602	2.947
16	1.337	1.746	2.120	2.583	2.921
17	1.333	1.740	2.110	2.567	2.898
18	1.330	1.734	2.101	2.552	2.878
19	1.328	1.729	2.093	2.539	2.861
20	1.325	1.725	2.086	2.528	2.845
21	1.323	1.721	2.080	2.518	2.831
22	1.321	1.717	2.074	2.508	2.819
23	1.319	1.714	2.069	2.500	2.807
24	1.318	1.711	2.064	2.492	2.797
25	1.316	1.708	2.060	2.485	2.787
26	1.315	1.706	2.056	2.479	2.779
27	1.314	1.703	2.052	2.473	2.771
28	1.313	1.701	2.048	2.467	2.763
29	1.311	1.699	2.045	2.462	2.756
30	1.310	1.697	2.042	2.457	2.750
31	1.309	1.696	2.040	2.453	2.744
32	1.309	1.694	2.037	2.449	2.738
33	1.308	1.692	2.035	2.445	2.733
34	1.307	1.691	2.032	2.441	2.728
35	1.306	1.690	2.030	2.438	2.724
36	1.306	1.688	2.028	2.434	2.719
37	1.305	1.687	2.026	2.431	2.715
38	1.304	1.686	2.024	2.429	2.712
39	1.304	1.685	2.023	2.426	2.708
40	1.303	1.684	2.021	2.423	2.704
50	1.299	1.676	2.009	2.403	2.678
60	1.296	1.671	2.000	2.390	2.660
70	1.294	1.667	1.994	2.381	2.646
80	1.292	1.664	1.990	2.374	2.639
90	1.291	1.662	1.987	2.368	2.632
100	1.290	1.660	1.984	2.364	2.626
110	1.289	1.659	1.982	2.361	2.621
120	1.289	1.658	1.980	2.358	2.617
∞	1.282	1.645	1.960	2.326	2.576

Source: This table was generated using the SAS® function TINV.