

1. (20%) The following eigen value problem is given as

$$\nabla^2 \Phi(x, y) = 0$$

in the domain $0 \leq x \leq \pi$ and $-h \leq y \leq 0$, where h is positive

constant, with the boundary conditions:

$$\frac{\partial \Phi}{\partial x} = 0 \quad \text{for } x = 0, -h \leq y \leq 0$$

$$x = \pi, -h \leq y \leq 0$$

$$\frac{\partial \Phi}{\partial y} = 0 \quad \text{for } 0 \leq x \leq \pi, y = -h$$

$$\frac{\partial \Phi}{\partial y} = \lambda \Phi \quad \text{for } 0 \leq x \leq \pi, y = 0.$$

- (1) (14%) Find all the eigen values λ_n and the corresponding eigen

functions Φ_n in terms of h .

- (2) (6%) Find λ_n in the limits $h \rightarrow 0$, and also $h \rightarrow \infty$.

2. (20%) For a function $f(z)$ is given as:

$$f(z) = \frac{1}{z^2 + z + 1} \ln\left(\frac{z-i}{z+i}\right)$$

- (1) (10%) Identify all singularities of $f(z)$, and specify their types.

- (2) (10%) Using $z-i = r_1 e^{i\theta_1}$, $z+i = r_2 e^{i\theta_2}$. Is $f(z)$ single valued?

Why and prove it.

3. (20%) Solve for $y(t)$ with $y(t) = u(t) + \int_0^t e^{-(t-\tau)} y(\tau) d\tau$ by using the

$$\text{Laplace Transform, where } u(t) = \begin{cases} 1 & t > 0 \\ 0 & t < 0 \end{cases}.$$

4. (a) (10%) Verify that $2x \frac{\partial F}{\partial x} - t \frac{\partial F}{\partial t} = 0$ has a solution $F(x,t) = f(x^\alpha t)$,

where f is arbitrary. Find the value of constant α .

(b) (10%) Solve this equation by separation of variables. By using superposition, obtain again the result above for arbitrary f . How is the proper α obtained in this approach?

5. (1) (10%) The nonlinear system

$$u_t + uu_x + gy_x = 0$$

$$y_t + uy_x + yu_x = 0 \quad \text{has the solution } u = u_0, y = y_0 \text{ (constants).}$$

Linearize the system by introducing the small perturbations:

$$u = u_0 + \varepsilon \omega(x,t), \quad y = y_0 + \varepsilon \eta(x,t) \quad \text{and neglect } \varepsilon^2 \text{ and higher order}$$

terms. Find this linearized system.

(2) (10%) Following (1) above by eliminating η from the system, find a second order linear differential equation for ω alone. Similarly, get an equation for η alone and compare the two equations.