

Numerical Analysis

1. (20%) Determine c_1 , c_2 , x_1 , and x_2 so that the integration formula

$$\int_{-1}^1 f(x) dx \approx c_1 f(x_1) + c_2 f(x_2)$$

gives the exact result whenever $f(x)$ is a polynomial of degree 3 or less.

2. (20%) Describe the Gauss-Seidel method for solving a linear system $Ax = b$, and find the first two iterations of this method for the system with

$$A = \begin{bmatrix} 3 & -1 & 1 \\ 3 & 6 & 2 \\ 3 & 3 & 7 \end{bmatrix} \quad b = \begin{bmatrix} 1 \\ 0 \\ 4 \end{bmatrix}$$

using $x^{(0)} = 0$.

3. (20%) Factor the matrix

$$A = \begin{bmatrix} 2 & -1 & 1 \\ 3 & 3 & 9 \\ 3 & 3 & 5 \end{bmatrix}$$

into the LU decomposition with $l_{ii} = 1$ for all i .

4. (20%) Write down a computer program in C or FORTRAN to apply Taylor's method of order 2 to solve the initial-value problem

$$y' = y - t^2 + 1, \quad 0 \leq t \leq 2, \quad y(0) = 0.5$$

5. (20%) Write an algorithm to apply the Newton's method for finding the solution of the nonlinear system

$$\begin{aligned} 5x_1^2 - x_2^2 &= 0 \\ x_2 - 0.25(\sin x_1 + \cos x_2) &= 0 \end{aligned}$$

using $(\frac{1}{4}, \frac{1}{4})^t$ as the starting point.