

1. Recall that for a sequence $\{x_n\}_1^{+\infty}$ in \mathbb{R} , $\liminf_{n \rightarrow +\infty} x_n$ is defined to be

$$\liminf_{n \rightarrow +\infty} x_n = \lim_{n \rightarrow +\infty} \left(\inf_{k \geq n} x_k \right)$$

Show that if $\lim_{n \rightarrow +\infty} y_n$ exists in \mathbb{R} , then for any sequence $\{x_n\}_1^{+\infty}$ in \mathbb{R} , one has
 $\liminf_{n \rightarrow +\infty} (x_n + y_n) = \liminf_{n \rightarrow +\infty} x_n + \lim_{n \rightarrow +\infty} y_n$. 15%

2. Suppose $\{f_n\}_1^{+\infty}$ is a sequence of continuous functions on $[0, +\infty)$. Suppose also there is a function g such that for any $n \in \mathbb{N}$, $|f_n(x)| \leq g(x)$ on $[0, +\infty)$ and $\int_0^{+\infty} g(x)dx$ converges. Prove that if $\{f_n\}_1^{+\infty} \rightarrow f$ uniformly on $[0, L]$ for any $L > 0$, then $\lim_{n \rightarrow +\infty} \int_0^{+\infty} f_n(x)dx = \int_0^{+\infty} f(x)dx$. 15%

3. (a) Suppose $f : [0, +\infty) \rightarrow [0, +\infty)$ is continuous and $\int_0^{+\infty} f(x)dx < +\infty$. Does it follow that $\lim_{x \rightarrow +\infty} f(x) = 0$? Support your statement with either a proof or a counterexample. 10%
- (b) If the continuity on f in (a) is replaced by uniform continuity, does it follow that $\lim_{x \rightarrow +\infty} f(x) = 0$? Support your statement with either a proof or a counterexample. 10%

4. Let $f(x, y)$ be a real-valued function on \mathbb{R}^2 which has continuous second partial derivatives. Show that f is harmonic if and only if

$$\int_C \left(\frac{\partial f}{\partial x} dy - \frac{\partial f}{\partial y} dx \right) = 0$$

for all circles C in the counterclockwise direction in \mathbb{R}^2 .

(Recall that f is harmonic if and only if $\frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} \equiv 0$ on \mathbb{R}^2 .) 15%

5. Let E be a region in \mathbb{R}^2 which is bounded by $3y - x = 0$, $3y - x + 1 = 0$, $y + 2x = 0$, and $y + 2x - 1 = 0$. Evaluate

$$\iint_E \sqrt[3]{2x^2 - 5xy - 3y^2} d(x, y). \quad 15\%$$

6. Let $f(x, y) = (x + y, \sin x + \cos y)$, $(x, y) \in \mathbb{R}^2$.
- (a) Show that a differentiable f^{-1} exists on some open set containing $(0, 1)$. 10%
- (b) Compute the matrix representation relative to the standard basis on \mathbb{R}^2 for the total derivative of f^{-1} at $(0, 1)$. 10%