

系所組別：統計學系學士班

考試科目：統計學

考試日期：0711，節次：4

※ 考生請注意：本試題 可 不可 使用計算機

1. Fifty male customers were asked which of two electric shavers, brand 1 or brand 2, they preferred. Nineteen of them preferred brand 1.
 - a. Find a 95% confidence interval for the proportion of male shoppers who preferred brand 1 over brand 2. (7 %)
 - b. Assume you have no prior knowledge of p (the proportion of males who preferred brand 1 over brand 2). Estimate the sample size that is required to estimate p , with 90% confidence, assuming a maximum error of .08. (8 %)

2. The following regression equation was computed from a sample of 20 observations: $\hat{Y} = 15 - 5X$. SSE was found to be 100 and SS total 400.
 - a. Determine the standard error of estimate. (5 %)
 - b. Determine the coefficient of correlation. (5 %)

3. A two-way factorial experiment was performed using two factors : type of tire and inflation level. The tires were driven 10,000 miles and the amount of tire wear was recorded. The inflation levels were slightly under inflated, correctly inflated, and slightly over inflated. Four replications were used for each combination of factor levels. The following ANOVA table for this design is partially filled in.

Source	df	SS	MS	F
Type of Tires	2		8.361	
Inflation				
Interaction			4.102	
Error		110.75		
Total		420.972		

- a. Complete the ANOVA table. (12 %)
 - b. Is there a significant difference in the levels of the type-of-tire factor? Use a 5% significance level. (6 %)
4. In an effort to monitor the service of its employees, a parcel-delivery firm keeps a tally of the number of packages misrouted each week at each of its 25 distribution centers, with the following results :

Misrouted Packages	0	1	2	3	>3
Distribution Centers	5	6	8	6	0

Do these data appear to come from a Poisson distribution with mean 1.6? Use a 0.05 significance level. (16 %)

(背面仍有題目,請繼續作答)

系所組別：統計學系學士班

考試科目：統計學

考試日期：0711，節次：4

※ 考生請注意：本試題 可 不可 使用計算機

5. An education testing service administers a national test to measure the reading comprehension of third-grade students. Only standardized scores are reported. The scores are approximately normally distributed. Find the z-scores corresponding to the 75th, 50th, and 20th percentiles.
(12 %)
6. A small company has 20% white-collar workers and 80% blue-collar workers. Of the white-collar workers, 50% believe that they would be financially secure for at least six months if they lost their job. Of the blue-collar workers, 30% believe that they would be financially secure for at least six months if they lost their job.
- What is the probability that a company employee believes that he/she would be financially secure if he/she lost his/her job? (6 %)
 - If a company employee believes that he/she would be financially secure for at least six months if he/she lost his/her job, what is the probability that the employee is a white-collar worker? (7 %)
7. Arnold Palmer and Tiger Woods are two of the best golfers to ever play the game. To show how these two golfers would compare if both were playing at the top of their game, the following sample data provide the results of 18-hole scores during a PGA tournament competition. Palmer's scores are from his 1960 season, while Wood's scores are from his 1999 season.

Palmer, 1960	Woods, 1999
$n_1 = 112$	$n_2 = 84$
$\bar{x}_1 = 69.95$	$\bar{x}_2 = 69.56$

Use the sample results to test the hypothesis of no difference between the population mean 18-hole scores for the two golfers.

- Assume a population variance of 6.25 for both golfers. What is the p -value? (13 %)
- At $\alpha = 0.01$, what is your conclusion? (3 %)


系所組別： 統計學系學士班

考試科目： 統計學

考試日期： 0711， 節次： 4

※ 考生請注意： 本試題 可 不可 使用計算機

TABLE Standard Normal Distribution



Entries in the table give the area under the curve between the mean and z standard deviations above the mean. For example, for $z = 1.25$ the area under the curve between the mean and z is .3944.

z	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
.0	.0000	.0040	.0080	.0120	.0160	.0199	.0239	.0279	.0319	.0359
.1	.0398	.0438	.0478	.0517	.0557	.0596	.0636	.0675	.0714	.0753
.2	.0793	.0832	.0871	.0910	.0948	.0987	.1026	.1064	.1103	.1141
.3	.1179	.1217	.1255	.1293	.1331	.1368	.1406	.1443	.1480	.1517
.4	.1554	.1591	.1628	.1664	.1700	.1736	.1772	.1808	.1844	.1879
.5	.1915	.1950	.1985	.2019	.2054	.2088	.2123	.2157	.2190	.2224
.6	.2257	.2291	.2324	.2357	.2389	.2422	.2454	.2486	.2518	.2549
.7	.2580	.2612	.2642	.2673	.2704	.2734	.2764	.2794	.2823	.2852
.8	.2881	.2910	.2939	.2967	.2995	.3023	.3051	.3078	.3106	.3133
.9	.3159	.3186	.3212	.3238	.3264	.3289	.3315	.3340	.3365	.3389
1.0	.3413	.3438	.3461	.3485	.3508	.3531	.3554	.3577	.3599	.3621
1.1	.3643	.3665	.3686	.3708	.3729	.3749	.3770	.3790	.3810	.3830
1.2	.3849	.3869	.3888	.3907	.3925	.3944	.3962	.3980	.3997	.4015
1.3	.4032	.4049	.4066	.4082	.4099	.4115	.4131	.4147	.4162	.4177
1.4	.4192	.4207	.4222	.4236	.4251	.4265	.4279	.4292	.4306	.4319
1.5	.4332	.4345	.4357	.4370	.4382	.4394	.4406	.4418	.4429	.4441
1.6	.4452	.4463	.4474	.4484	.4495	.4505	.4515	.4525	.4535	.4545
1.7	.4554	.4564	.4573	.4582	.4591	.4599	.4608	.4616	.4625	.4633
1.8	.4641	.4649	.4656	.4664	.4671	.4678	.4686	.4693	.4699	.4706
1.9	.4713	.4719	.4726	.4732	.4738	.4744	.4750	.4756	.4761	.4767
2.0	.4772	.4778	.4783	.4788	.4793	.4798	.4803	.4808	.4812	.4817
2.1	.4821	.4826	.4830	.4834	.4838	.4842	.4846	.4850	.4854	.4857
2.2	.4861	.4864	.4868	.4871	.4875	.4878	.4881	.4884	.4887	.4890
2.3	.4893	.4896	.4898	.4901	.4904	.4906	.4909	.4911	.4913	.4916
2.4	.4918	.4920	.4922	.4925	.4927	.4929	.4931	.4932	.4934	.4936
2.5	.4938	.4940	.4941	.4943	.4945	.4946	.4948	.4949	.4951	.4952
2.6	.4953	.4955	.4956	.4957	.4959	.4960	.4961	.4962	.4963	.4964
2.7	.4965	.4966	.4967	.4968	.4969	.4970	.4971	.4972	.4973	.4974
2.8	.4974	.4975	.4976	.4977	.4977	.4978	.4979	.4979	.4980	.4981
2.9	.4981	.4982	.4982	.4983	.4984	.4984	.4985	.4985	.4986	.4986
3.0	.4986	.4987	.4987	.4988	.4988	.4989	.4989	.4989	.4990	.4990

(背面仍有題目,請繼續作答)


系所組別： 統計學系學士班

考試科目： 統計學

考試日期： 0711，節次： 4


※ 考生請注意：本試題 可 不可 使用計算機

Chi-Square Distribution Table



The shaded area is equal to α for $\chi^2 = \chi^2_{\alpha}$.

df	$\chi^2_{.995}$	$\chi^2_{.990}$	$\chi^2_{.975}$	$\chi^2_{.950}$	$\chi^2_{.900}$	$\chi^2_{.100}$	$\chi^2_{.050}$	$\chi^2_{.025}$	$\chi^2_{.010}$	$\chi^2_{.005}$
1	0.000	0.000	0.001	0.004	0.016	2.706	3.841	5.024	6.635	7.879
2	0.010	0.020	0.051	0.103	0.211	4.605	5.991	7.378	9.210	10.597
3	0.072	0.115	0.216	0.352	0.584	6.251	7.815	9.348	11.345	12.838
4	0.207	0.297	0.484	0.711	1.064	7.779	9.488	11.143	13.277	14.860
5	0.412	0.554	0.831	1.145	1.610	9.236	11.070	12.833	15.086	16.750
6	0.676	0.872	1.237	1.635	2.204	10.645	12.592	14.449	16.812	18.548
7	0.989	1.239	1.690	2.167	2.833	12.017	14.067	16.013	18.475	20.278
8	1.344	1.646	2.180	2.733	3.490	13.362	15.507	17.535	20.090	21.955
9	1.735	2.088	2.700	3.325	4.168	14.684	16.919	19.023	21.666	23.589
10	2.156	2.558	3.247	3.940	4.865	15.987	18.307	20.483	23.209	25.188



TABLE

Critical Values of F

For a particular combination of numerator and denominator degrees of freedom, entry represents the critical values of F corresponding to the cumulative probability $(1 - \alpha)$ and a specified upper-tail area (α) .

Denominator, df_2	Numerator, df_1													
	1	2	3	4	5	6	7	8	9	10	12	15	30	40
1	161.40	199.50	215.70	224.60	230.20	234.00	236.80	238.90	240.50	241.90	243.90	245.90	250.10	251.10
2	18.51	19.00	19.16	19.25	19.30	19.33	19.35	19.37	19.38	19.40	19.41	19.43	19.46	19.47
3	10.13	9.55	9.28	9.12	9.01	8.94	8.89	8.85	8.81	8.79	8.74	8.70	8.62	8.59
4	7.71	6.94	6.59	6.39	6.26	6.16	6.09	6.04	6.00	5.96	5.91	5.86	5.75	5.72
23	4.28	3.42	3.03	2.80	2.64	2.53	2.44	2.37	2.32	2.27	2.25	2.22	2.16	2.14
24	4.26	3.40	3.01	2.78	2.62	2.51	2.42	2.36	2.30	2.25	2.18	2.11	1.94	1.91
25	4.24	3.39	2.99	2.76	2.60	2.49	2.40	2.34	2.28	2.24	2.16	2.09	1.92	1.87
26	4.23	3.37	2.98	2.74	2.59	2.47	2.39	2.32	2.27	2.22	2.15	2.07	1.90	1.85
27	4.21	3.35	2.96	2.73	2.57	2.46	2.37	2.31	2.25	2.20	2.13	2.06	1.88	1.84
28	4.20	3.34	2.95	2.71	2.56	2.45	2.36	2.29	2.24	2.19	2.12	2.04	1.87	1.82
29	4.18	3.33	2.93	2.70	2.55	2.43	2.35	2.28	2.22	2.18	2.10	2.03	1.85	1.81
30	4.17	3.32	2.92	2.69	2.53	2.42	2.33	2.27	2.21	2.16	2.09	2.01	1.84	1.79
40	4.08	3.23	2.84	2.61	2.45	2.34	2.25	2.18	2.12	2.08	2.00	1.92	1.74	1.69
60	4.00	3.15	2.76	2.53	2.37	2.25	2.17	2.10	2.04	1.99	1.92	1.84	1.65	1.59
120	3.92	3.07	2.68	2.45	2.29	2.17	2.09	2.02	1.96	1.91	1.83	1.75	1.55	1.50
∞	3.84	3.00	2.60	2.37	2.21	2.10	2.01	1.94	1.88	1.83	1.75	1.67	1.46	1.39